



#6

SEQUENCE LISTING

<110> INOUE, Akio
SHIMIZU, Yoshihiro
UEDA, Takuya

<120> Process for Producing Peptides By Using In Vitro
Transcription/Translation System

<130> 1752-0151P

<140> 09/983,067

<141> 2001-10-23

<150> JP 294795/2001

<151> 2001-09-26

<150> JP 227094/2001

<151> 2001-07-27

<150> JP 6910/2001

<151> 2001-01-15

<150> JP 401417/2000

<151> 2000-12-28

<160> 6

<170> PatentIn version 3.1

<210> 1

<211> 14

<212> PRT

<213> Bovine sp.

<400> 1

Ser Asn Lys Glu Gln Glu Pro Lys Glu Lys Lys Lys Lys Lys
1 5 10

<210> 2

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Strep-tag binding to streptavidin

<220>

<221> MISC_FEATURE

<222> (1)..(9)

<223> Schmidt & Skerra, 1993, "The random peptide library-assisted
enginerring of a C-terminal affinity peptide, useful for the
detection and purification of a functional IgFv fragment",
Protein Eng. 6(1):109-122.

<400> 2

Ala Trp Arg His Pro Gln Phe Gly Gly
1 5

<210> 3

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Strep-tag II used in the binding of streptavidin

<220>

<221> MISC_FEATURE

<222> (1)..(9)

<223> Schmidt et al., 1996, "Molecular interaction between the
Strep-tag affinity peptide and its cognate target,
streptavidin", J. of Mol. Biol. 255(5):753-766.

<400> 3

Asn Trp Ser His Pro Gln Phe Glu Lys
1 5

<210> 4

<211> 12

<212> RNA

<213> Artificial Sequence

<220>

<223> RNA sequence to obtain a template for MFL mRNA

<220>

<221> mRNA

<222> (1)..(12)

<223>

<400> 4

auguucuugu aa

12

<210> 5

<211> 14

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide A that anneals to SEQ ID NO:6 to give a
double-stranded DNA containing NdeI and XhoI sequences

<400> 5
tatgttcttg taac

14

<210> 6
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide B that anneals to SEQ ID NO: 5 to give a
double-stranded DNA containing NdeI and XhoI sequences

<400> 6
tcgagttaca agaaca

16